



## **OrgTrace – No difference found in bioactive compounds of organic and conventional crops**

**Knuthsen, Pia; Søltoft, Malene; Laursen, Kristian Holst; Bysted, Anette; Madsen, Katja Howarth; Nielsen, John; Halekoh, Ulrich; Husted, Søren**

*Publication date:*  
2011

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Knuthsen, P., Søltoft, M., Laursen, K. H., Bysted, A., Madsen, K. H., Nielsen, J., Halekoh, U., & Husted, S. (2011). *OrgTrace – No difference found in bioactive compounds of organic and conventional crops*. Poster session presented at 1st International Conference on Organic Food Quality and Health Research, Prague, Czech Republic.

---

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



# OrgTrace – No difference found in bioactive compounds of organic and conventional crops

P. Knuthsen<sup>1</sup>\*, M. Søltoft<sup>1</sup>, K.H. Laursen<sup>2</sup>, A. Bysted<sup>1</sup>, K.H. Madsen<sup>1</sup>, J. Nielsen<sup>2</sup>, U. Halekoh<sup>3</sup>, and S. Husted<sup>2</sup>

<sup>1</sup> DTU Food, National Food Institute, Technical University of Denmark, DK-2860 Søborg

<sup>2</sup> Faculty of Life Sciences, University of Copenhagen, DK-1871, Denmark

<sup>3</sup> Faculty of Agricultural Sciences, Aarhus University, DK-8830 Tjele, Denmark

## Introduction

Secondary plant metabolites, like carotenoids, flavonoids, phenolic acids and polyacetylenes, are bioactive compounds with presumably beneficial health effects. Contents in plants are affected by e.g. Plant nutrient availability, climate, and pathogen infection.

## Objectives


The objectives of our study was to compare the content of selected carotenoids, flavonoids, phenolic acids, and polyacetylenes in organically and conventionally grown carrots, onions, and potatoes. And to evaluate if the ability of the crops to synthesize selected secondary metabolites is systematically affected by growth systems across different growth years as well as geographic locations (soil types).

## Study design

Carrots, onions, and potatoes were cultivated in two-year field trials in three different geographical locations, comprising one conventional (C) and two organic agricultural systems (OA relying on import of animal manure, and OB based on the use of cover crops). The crops were harvested at the same day for all systems, and representative sampling ensured.



PLE (Pressurized Liquid Extraction) - for extraction of flavonoids and phenolic acids before analysis.



Carotenoids in carrots.  
Lutein – α-carotene – β-carotene  
Year 1 and year 2.

## Analyses

The edible parts of the crops were freeze-dried for analysis, homogenized, and stored at 20 °C in nitrogen atmosphere, protected from light and oxygen during entire sample preparation. The bioactive compounds were extracted by optimized methods, and further identified and quantified by LC-MS-MS analyses.

## Results

No systematic effects of growth system, year or location was found on contents of the selected secondary metabolites - carotenoids, flavonoids, phenolic acids, and polyacetylenes - in the carrots, onions or potatoes.

## References:

- M. Søltoft et al. J. Agric. Food Chem. 2010, 58, 7673-7679
- M. Søltoft et al. J. Agric. Food Chem. 2010, 58, 10323-10329
- M. Søltoft et al. J. Sci. Food Agric. 2011, 91, 767-775